(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau





(43) International Publication Date 7 November 2002 (07.11.2002)

PCT

(10) International Publication Number WO 02/088342 A1

- (51) International Patent Classification7: C12N 9/00, 9/98, 1/14, C12P 21/00, A21D 8/04 // (C12N 9/00, C12R 1:685) (C12N 9/00, C12R 1:845) (C12N 9/98, C12R 1:685) (C12N 9/98, C12R 1:845) (C12P 21/00, C12R 1:685) (C12P 21/00, C12R 1:845)
- (21) International Application Number: PCT/IN01/00094
- (22) International Filing Date: 30 April 2001 (30.04.2001)
- (25) Filing Language:

English

(26) Publication Language:

English

- (71) Applicant (for all designated States except US): BIOCON INDIA LIMITED [IN/IN]; 20th Km Hosur Road, Hebbagodi, Bangalore 561 229, Karnataka (IN).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): RANGANATHAN, Lakshmi [IN/IN]; 20th Km Hosur Road, Hebbagodi, Bangalore 561 229, Karnataka (IN). JAYARAM, Shiv, Kumar [IN/IN]; 20th Km Hosur Road, Hebbagodi, Bangalore 561 229, Karnataka (IN). KAMATH, Jyothi, Anantha [IN/IN]; 20th Km Hosur Road, Hebbagodi, Bangalore 561 229, Karnataka (IN).

- (74) Agents: ANAND, Pravin et al.; Anand & Anand Advocates, B-41, Nizamuddin East, New Delhi 110 013 (IN).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

- of inventorship (Rule 4.17(iv)) for US only

Published:

- with international search report
- with amended claims

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: AN ENZYME PREPARATION FOR IMPROVED BAKING QUALITY AND A PROCESS FOR PREPARING THE SAME

(57) Abstract: The present invention provides an improved method for the preparation of an enzyme mixture for improved fresh-keeping of baked products. The enzyme preparation is produced by the co-culture of two fungal species of *Rhizopus & Aspergillus* by solid substrate fermentation under optimal fermentation parameters. The product is extracted and further processed to get an enzyme preparation as solid powder. The present invention also includes an enzyme preparation obtained from *Rhizopus sp.* and *Aspergillus sp.*.



AN ENZYME PREPARATION FOR IMPROVED BAKING QUALITY AND A PROCESS FOR PREPARING THE SAME

The present invention relates to an enzyme preparation for improved baking quality and a process for preparing the same

BACKGROUND

The process of baking has been in existence for at least 6000 years and has been exploited ever since to improve the process. The major components of baked products are polysaccharides, proteins, fats & emulsifiers, enzymes, yeast and water. Of these enzymes are involved in the oxidative aging of flour, in dough development and in the formation of bread flavor & texture. They are also responsible for a number of adverse effects on the quality of the baked product. On the whole the overall effect of the indigenous enzymes of the grain, the microorganisms & of various added exogenous enzymes influence the quality of the baked product (Chemistry and physics of baking. Ed. J.M.V. Blanshard, P. J. Frazier & T. Galliard, Published by The Royal Society of Chemistry 1986).

Staling of baked products is a major negative influence & is defined as an increase of crumb firmness & a corresponding loss of moisture & freshness. During staling there is a gradual transition of the polysaccharide starch from an amorphous to a partially crystalline state. This increase in crystallinity called retrogradation is due to an inter or intra molecular association of starch molecules via hydrogen bonding. The staling process is influenced by time, temperature, moisture & presence of additives in the baked product (Enzymes used in the milling industry. Ter Haseborg .E Alimenta, 1988, 27, 1, 2-10).

1

Additives act by way of complexing with the components of starch - amylose & amylopectin, thereby reducing their tendency to retrograde. Increasingly, enzymes too are being added to prevent staling. These antistaling enzymes are generally alpha & beta amylases, glucoamylases, glucose oxidase, peroxidases and lipoxygenases.

For a successful antistaling enzyme it is essential that they can act on raw starch rather than gelatinised starch & have an intermediate heat stability i.e. more stable than the conventional heat labile fungal amylases but less stable than the conventional heat stable bacterial amylases. These enzymes would thus provide an antistaling effect without adversely affecting the quality of the baked product, whereby they hydrolyse starch during the dough development process, but are completely heat inactivated before the baking process is completed. They thus prevent an excess of the breakdown products of starch—the dextrins which lead to an increased stickiness & gumminess of the bread (Enzymes for the baking industry. Van Oort, M.G., Hamer, R.J. Aliment., Equipos Tecnol, 1993, 12,5,115–18). The antistaling enzyme should, besides aiding fresh keeping of bread should not adversely affect the other desirable parameters of a baked product like loaf volume, color, softness, resilience & crumb texture (Synergistic effect of enzymes for breadbaking. Si, J. Qi Cereal Foods World 9, 1997, 42, 10,802-807).

Co culture of different strains of bacteria - bacteria (Deanda et. al., Appl Environ Microbiol, 1996,62,12,4465-70), yeast -bacteria (Kim et. al., Biotechnol Lett, 1996,18,9,1031-34), yeast -yeast , yeast / bacteria - fungi (Padmaja et. al. J. Sci Food Agric, 1993, 63,4,473-81; Marakis S.G Biotechnol Lett, 1992,14,11,1075-80) & fungi -fungi (Benzuela Elegado, Francisco, Fujio, Yusaku J. Gen. Appl.

Microbiol. 1993, 39,6, 541-6; Gutierrez-Correa, Marcel; Tengerdy, Robert P, Biotechnol. Lett. 1997,19,7,665-667; Cellulase production by mixed fungi in solid substrate fermentation of bagasse (Duenas R et. Al. World T. Microbiol. Biotechnol ,1995 ,11,3,333-337) have been reported in literature. These associations have been used to produce many specialty products like acetic acid, alcohol butanol & enzymes like xylanase (Xylanase production by fungal mixed culture solid substrate fermentation on sugarcane bagasse. Gutierrez-Correa, Marcel; Tengerdy, Robert P, Biotechnol. Lett. 1998, 20, 1, 45 - 47) and cellulases (Denada et. al., Appl Environ Microbiol, 1996,62 12,4465-70). The latter has been obtained using Trichoderma & Aspergillus strains (Mixed culture solid substrate fermentation of Trichoderma reesei with Aspergillus niger on sugarcane bagasse. Gutierrez-Correa, Marcel; Portal Leticia, Moreno Patricia, Tengerdy, Robert P. Bioresour. Technol, 1998, Volume date 1999, 68,2,173-178; Production of cellulase on sugarcane bagasse by fungal mixed culture solid substrate fermentation. Gutierrez-Correa, Marcel; Tengerdy, Robert P, Biotechnol. Lett. 1997,19,7,665-667; Formation of cellulases by co-culturing of Trichoderma reesei and Aspergillus niger on cellulosic waste. Madamwar, D, Patel, S, World J. Microbiol.Biotechnol 1992, 8,2,183-6).

DESCRIPTION OF THE INVENTION

The present invention relates to a novel process to produce an enzyme preparation with antistaling properties for improved freshkeeping of baked products.

Accordingly, the present invention provides for a A method for the manufacture of an enzyme preparation for improved baked products, which comprises of:

a. preparing an inoculum of Rhizopus sp.,

- b. preparing an inoculum of Aspergillus sp.,
- c. mixing the grown inoculum of Rhizopus sp. and Aspergillus sp.,
- d. sterilizing a solid state nutritive matrix,
- e. mixing the said sterilized solid state nutritive matrix with the mixture of inoculum consisting of *Rhizopus sp.* and *Aspergillus sp.*,
- f. incubating the said inoculated solid state nutritive matrix for 4 7 days at 25 35°C,
- g. extracting the fermented matrix followed by filtration,
- h. concentrating the aqueous extract,
- i. spray drying of the concentrated aqueous extract to get the **enzyme** preparation.

The Rhizopus sp. is Rhizopus oryzae. The Aspergillus sp. is Aspergillus niger. The inoculum of Rhizopus sp. and Aspergillus sp. is mixed in the ratio ranging from 3:97 to 97:3. The inoculum of Rhizopus sp. and Aspergillus sp. is mixed in the ratio of 12.5:87.5. The solid state nutritive matrix is selected from wheat bran, rice bran, soya grits, rice grits, millet flour, soya flour, sugar beet, bagasse or a mixture of these. The mixed inoculum is optionally grown prior to mixing with sterilized solid state matrix. The extraction is carried out using water or an aqueous buffer. The aqueous extract is concentrated by ultra-filtration or reverse osmosis. In step (e) at least 10 % of inoculum is mixed with the solid state matrix. In step (f), the inoculate solid state matrix is incubated at 30 °C. The quantity of water added to the extract is in the ration of 1:6. The spray dried enzyme preparation is granulated. The spray dried enzyme preparation contains a preservative. The preservative is selected from benzoates or sorbates. The spray dried enzyme preparation contains a stabilizer. The stabilizer is selected from inorganic salts, polyols, sugars or their combinations. The spray drying step in 1 (i) is replaced by freeze drying or solvent precipitation. The inoculum is optionally not mixed but added together to the sterilized nutritive matrix.

The enzyme preparation is used for baking for improved freshkeeping, whitening, softening, crumb texture or volume increase.

The present invention also comprises an enzyme preparation for improving baked products obtained from *Rhizopus sp.* and *Aspergillus sp.* The *Rhizopus sp.* is *Rhizopus oryzae*. The *Aspergillus sp.* is *Aspergillus niger*. The enzyme preparation further comprises a preservative, preferably selected from benzoates or sorbates. The enzyme preparation further comprises a stabilizer, preferably selected from inorganic salts, polyols, sugars or their combinations.

The present invention has the following advantages over the other reported methods:

- (i) The enzyme preparation obtained is from a non-genetically modified (GMO) origin.
- (ii) The process is environment friendly, less cumbersome and economical for large-scale industrial applications.
- (iii) A broader range of enzyme activities that aid freshkeeping is obtained, which is not obtained with a single culture alone.
- (iv) This combination of desired activities for freshkeeping is generated *in situ* thereby having an advantage over blended enzyme preparations.
- (v) The enzyme preparation is capable of acting on raw starch, rather than gelatinized starch.

The invention will now be described with reference to the following examples:

EXAMPLE 1

Preparation of inoculum of Rhizopus sp:

About 100 µL spore suspension of *Rhizopus oryzae*, made by adding 3 mL of sterile water to a culture slant, is added to a 250 mL Erlenmeyer flask containing 35 mL of medium with following composition:

No	Ingredient	Quantity	
		(%)	
1	Wheat flour	4.14	
2	Sucrose	1	
3	Peptone	0.306	
4	Ammonium sulphate	0.2	
5	Yeast extract	0.2	
6	Potassium dihydrogen phosphate	0.085	
7	CaCl ₂ .2H ₂ O	0.01	
8	MgSO ₄ . 7 H ₂ O	0.01	
9	NaCl	0.01	

The flasks are incubated at 30 deg C on a rotary shaker (200-rpm) for 48 hours.

EXAMPLE 2

Preparation of inoculum of Aspergillus sp:

About 100 µL spore suspension of *Aspergillus niger*, made by adding 3 mL of sterile water to a culture slant, is added to a 250 mL Erlenmeyer flask containing 35 mL of medium with following composition:

No	Ingredient	Quantity
		(%)
1	Wheat flour	4.14

2	Peptone	0.306
3	Mono ammonium phosphate	0.0162
4	Amycoglucosidase	0.0054

The flasks are incubated at 30 deg C on a rotary shaker (200-rpm) for 48 hours.

EXAMPLE 3

Mixing of the inoculum

The inoculum developed from Example 1 is mixed the inoculum of Example 2 in the ratio 12.5:87.5. This mixed inoculum is used in the subsequent stages.

Example 4

Sterilization of wheat bran:

Commercial wheat bran is used as the solid substrate. Moist wheat bran is autoclaved for 90minutes in a rotating cooker.

Example 5

Inoculation:

The mixed inoculum from example 3 is mixed with the sterilized wheat bran from example 4. A 10 % inoculum is used. The inoculum is mixed at 30 deg. C. After a proper mixing, the inoculated wheat bran is transferred and layered into the presterilized trays.

Example 6

Incubation:

Incubation of the layered trays from example 5 is done at 30 deg.C. for 5 days. At the end of 5 days, the fermented solid substrate is harvested.

Example 7

Extraction:

Extraction is done using water. Quantity of water added to the extract is in a ratio of 1:6. Soaking of the harvested solid substrate is done for 6 hours.

Example 8

Extraction:

The extraction is carried out in the same way as example 7 but with the addition of 0.2% sodium benzoate with the water.

Example 9

Microfiltration:

Microfiltartion is done to the extract obtained from example 7 to obtain a microorganism free product. This is done using a 0.2 micron MF membrane.

Example 10

Ultrafiltration:

The resultant liquid obtained from example 9 is concentrated to the desired activity by Ultrafiltration done at 15 deg.C. Ultrafiltration is done using UF membranes.

Example 11

Enzyme stabilization:

The Ultrafiltered enzyme obtained from example 10 is stabilized by the addition of NaCl and KCl.

Example 12

Enzyme stabilization:

The Ultrafiltered enzyme obtained from example 10 is stabilized by the addition of gylcerol and sorbitol.

Example 13

Spray drying:

The UFC (ultrafiltered concentrate) as obtained from example 10 is spray dried to get the required enzyme preparation.

Example 14

Spray drying:

The UFC (ultrafiltered concentrate) as obtained from Example 11 is spray dried as in Example 13, but with the addition of 0.1 % potassium sorbate as a preservative before spray drying.

Example 15

Freeze drying:

The UFC (ultrafiltered concentrate) as obtained from Example 11 is freeze dried.

Example 16

Baking trials:

9

The enzyme preparation obtained from example 15 was used in a baking trial. This freeze dried enzyme was dosed at 113 ppm in a bread baking trial & was evaluated for fresh keeping against a trial which did not have the enzyme.

The results of the baking trials of the test sample (containing the enzyme preparation), were compared with the control (where the enzyme preparation was absent) and the quality of the bread evaluated. The following observations were made:

DAYS	EVALUATION	CONTROL	TEST
	Enzyme	Absent	Present
	preparation		·
05	Softness	+	+++
	Mold Growth	-	-
	Slice strength	Good	Good
10	Softness	+	+++
	Mold Growth	+	-
	Slice strength	Weak	Weak
14	Softness	+	1-1-1
	Mold Growth	++	+
	Slice strength	Weak	Weak
	OVERALL	+	+++
	RATING		·

The bread made with the freeze dried enzyme preparation appeared softer than that of the control even after 14 days.

Note:

- + denotes poor
- ++ denotes average
- +++ denotes good
- denotes absent

We claim:

1. A method for the manufacture of an enzyme preparation for improved baked products, which comprises of:

- j. preparing an inoculum of Rhizopus sp.,
- k. preparing an inoculum of Aspergillus sp.,
- 1. mixing the grown inoculum of Rhizopus sp. and Aspergillus sp.,
- m. sterilizing a solid state nutritive matrix,
- n. mixing the said sterilized solid state nutritive matrix with the mixture of inoculum consisting of *Rhizopus sp.* and *Aspergillus sp.*,
- incubating the said inoculated solid state nutritive matrix for 4 7 days at 25 - 35°C,
- p. extracting the fermented matrix followed by filtration,
- q. concentrating the aqueous extract,
- r. spray drying of the concentrated aqueous extract to get the **enzyme** preparation.
- 2. The method as claimed in claim 1 wherein the Rhizopus sp. is Rhizopus oryzae.
- 3. The method as claimed in claim 1 wherein the Aspergillus sp. is Aspergillus niger.
- 4. The method as claimed in claim 1 wherein the inoculum of *Rhizopus sp.* and *Aspergillus sp.* is mixed in the ratio ranging from 3:97 to 97:3.
- 5. The method as claimed in claim 4 wherein the inoculum of *Rhizopus sp.* and *Aspergillus sp.* is mixed in the ratio of 12.5: 87.5.

6. The method as claimed in claim 1 wherein the solid state nutritive matrix is selected from wheat bran, rice bran, soya grits, rice grits, millet flour, soya flour, sugar beet, bagasse or a mixture of these.

- 7. The method as claimed in claim 1 wherein the mixed inoculum is optionally grown prior to mixing with sterilized solid state matrix.
- 8. The method as claimed in claim 1 wherein the extraction is carried out using water or an aqueous buffer.
- 9. The method as claimed in claim 1 wherein the aqueous extract is concentrated by ultra-filtration or reverse osmosis.
- 10. The method as claimed in claim 1 wherein in step (e) at least 10 % of inoculum is mixed with the solid state matrix.
- 11. The method as claimed in claim 1 wherein in step (f), the inoculate solid state matrix is incubated at 30 °C.
- 12. The method as claimed in claim 8 wherein quantity of water added to the extract is in the ration of 1:6.
- 13. The method as claimed in claim 1 wherein the spray dried enzyme preparation is granulated.
- 14. The method as claimed in claim 1 or 13 wherein the spray dried enzyme preparation contains a preservative.
- 15. The method as claimed in claim 14 wherein the preservative is selected from benzoates or sorbates.

The method as claimed in any one of the preceding claims wherein the spray dried enzyme preparation contains a stabilizer.

- 17. The method as claimed in claim 16 wherein the stabilizer is selected from inorganic salts, polyols, sugars or their combinations.
- 18. The method as claimed in claim 1 to 17 wherein the spray drying step in 1 (i) is replaced by freeze drying or solvent precipitation.
- 19. The method as claimed in claim 1 to 14 wherein the inoculum is optionally not mixed but added together to the sterilized nutritive matrix.
- 20. The method as claimed in claim 1 to 14 wherein the enzyme preparation is used for baking for improved freshkeeping, whitening, softening, crumb texture or volume increase.
- 21. An enzyme preparation for improving baked products obtained from *Rhizopus* sp. and *Aspergillus sp*.
- 22. An enzyme preparation as claimed in claim 21 wherein *Rhizopus sp.* is *Rhizopus oryzae*.
- 23. An enzyme preparation as claimed in claim 22 wherein the Aspergillus sp. is Aspergillus niger.
- 24. An enzyme preparation as claimed in claim 21 wherein the enzyme preparation is granulated.
- 25. An enzyme preparation as claimed in any one of claims 21- 24 further comprising a preservative.

26. An enzyme preparation as claimed in claim 25 wherein the preservative is selected from benzoates or sorbates.

- 27. An enzyme preparation as claimed in any one of the preceding claims further comprising a stabilizer.
- 28. An enzyme preparation as claimed in claim 27 wherein the stabilizer is selected from inorganic salts, polyols, sugars or their combinations.

5

10

15

20

AMENDED CLAIMS

[received by the International Bureau on 15 March 2002 (15.03.02); original claims 1-28 replaced by new claims 1-20 (3 pages)]

- 1. A method for the manufacture of an enzyme preparation for improved baked products, which comprises of:
 - a. preparing an inoculum of Rhizopus sp.,
 - b. preparing an inoculum of Aspergillus sp.,
 - c. mixing the grown inoculum of Rhizopus sp. and Aspergillus sp.,
 - d. sterilizing a solid state nutritive matrix,
 - e. mixing the said sterilized solid state nutritive matrix with the mixture of inoculum consisting of *Rhizopus sp.* and *Aspergillus sp.*,
 - f. incubating the said inoculated solid state nutritive matrix for 4 7 days at 25 35°C,
 - g. extracting the fermented matrix followed by filtration,
 - h. concentrating the aqueous extract,
 - i. spray drying of the concentrated aqueous extract to get the enzyme preparation.
- 2. The method as claimed in claim 1 wherein the Rhizopus sp. is Rhizopus oryzae.
- 3. The method as claimed in claim I wherein the Aspergillus sp. is Aspergillus niger.
- 4. The method as claimed in claim 1 wherein the inoculum of *Rhizopus sp.* and Aspergillus sp. is mixed in the ratio ranging from 3:97 to 97:3.
 - 5. The method as claimed in claim 4 wherein the inoculum of Rhizopus sp. and Aspergillus sp. is mixed in the ratio of 12.5: 87.5.

6. The method as claimed in claim 1 wherein the solid state nutritive matrix is selected from wheat bran, rice bran, soya grits, rice grits, millet flour, soya flour, sugar beet, bagasse or a mixture of these.

7. The method as claimed in claim 1 wherein the mixed inoculum is optionally grown prior to mixing with sterilized solid state matrix.

5

- 8. The method as claimed in claim I wherein the extraction is carried out using water or an aqueous buffer.
- 9. The method as claimed in claim 1 wherein the aqueous extract is concentrated by ultra-filtration or reverse osmosis.
- 10. The method as claimed in claim 1 wherein in step (e) at least 10 % of inoculum is mixed with the solid state matrix.
 - 11. The method as claimed in claim 1 wherein in step (f), the inoculate solid state matrix is incubated at 30 °C.
- 12. The method as claimed in claim 8 wherein quantity of water added to the extract is in the ration of 1:6.
 - 13. The method as claimed in claim 1 wherein the spray dried enzyme preparation is granulated.
 - 14. The method as claimed in claim 1 or 13 wherein the spray dried enzyme preparation contains a preservative.
- 20 15. The method as claimed in claim 14 wherein the preservative is selected from benzoates or sorbates.

16. The method as claimed in any one of the preceding claims wherein the spray dried enzyme preparation contains a stabilizer.

- 17. The method as claimed in claim 16 wherein the stabilizer is selected from inorganic salts, polyols, sugars or their combinations.
- 18. The method as claimed in claim 1 to 17 wherein the spray drying step in 1 (i) is replaced by freeze drying or solvent precipitation.
 - 19. The method as claimed in claim 1 to 14 wherein the inoculum is optionally not mixed but added together to the sterilized nutritive matrix.
- 20. The method as claimed in claim 1 to 14 wherein the enzyme preparation is used for baking for improved fresh keeping, whitening, softening, crumb texture or volume increase.

1.5

Inte ial Application No PC 1/1N 01/00094

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C12N9/00 C12N9/98 C12N1/14 C12P21/00 A21D8/04 //(C12N9/00,C12R1:685),(C12N9/00,C12R1:845),(C12N9/98,C12R1:685), (C12N9/98,C12R1:845),(C12P21/00,C12R1:685),(C12P21/00,C12R1:845)					
According to	According to International Patent Classification (IPC) or to both national classification and IPC				
	SEARCHED				
	Minimum documentation searched (classification system followed by classification symbols)				
	ion searched other than minimum documentation to the extent that su				
Electronic d	ata base consulted during the international search (name of data bas	e and, where practical, search terms used)		
	ternal, WPI Data, PAJ, BIOSIS, FSTA,	MEDLINE, EMBASE			
C. DOCUME	ENTS CONSIDERED TO BE RELEVANT				
Category *	Citation of document, with indication, where appropriate, of the rele	evant passages	Relevant to claim No.		
X	KIM H-S ET AL.: "Enzymological characteristics and identificatio useful fungi isolated from tradit Korean Nuruk." KOREAN JOURNAL OF APPLIED MICROBI BIOTECHNOLOGY, vol. 26, no. 5, 1998, pages 456-4 XP001051554 abstract table 8	ional OLOGY AND	1,2		
Further documents are listed in the continuation of box C. X Patent family members are listed in annex.					
 Special categories of cited documents: 'A' document defining the general state of the art which is not considered to be of particular relevance 'E' earlier document but published on or after the international filling date 'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) 'O' document referring to an oral disclosure, use, exhibition or other means 'P' document published prior to the international filling date but later than the priority date claimed Date of the actual completion of the international search 'I' later document published after the international filling date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken along the cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. 'A' document member of the same patent family Date of the actual completion of the international search 		the application but ecory underlying the claimed invention to economic the considered to cournent is taken alone claimed invention eventive step when the one other such docuus to a person skilled family			
	January 2002	23/01/2002			
Name and a	mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL – 2280 HV Rijswijk Tel. (+31–70) 340–2040, Tx. 31 651 epo nl,	Authorized officer Van de Kamp, M			
I	Fax: (+31-76) 340-3016				

Inte nal Application No PCT/IN 01/00094

to claim No.
,21
1,22, 5-28
4
21,23
24
.,21
· .
l
l

Inte 1 1 Application No
PC., N 01/00094

•	PC., IN 01/00094	
C.(Continua	ntion) DOCUMENTS CONSIDERED TO BE RELEVANT .	
Category °	senessen treveler of the relevant passages	Relevant to claim No.
A	BIGELIS R: "Handbook of applied mycology, Volume 3: Foods and feeds (eds Arora D K et al.), Chapter 14: FUNGAL ENZYMES IN FOOD PROCESSING (pp 445-498)" 1991 , MARCEL DEKKER, INC. , NEW YORK, USA; ISBN 0-8247-8491-X. XP001051671 page 445, line 15 -page 449, line 8 page 476, line 1 -page 478, line 9	
А	BOWLES L K: "Amylolytic enzymes. In: Baked goods freshness. Technology, evaluation, and inhibition of staling (pp 105-129)." 1996, MARCEL DEKKER, INC., NEW YORK, USA XP002186664 page 126, line 28 -page 127, line 12	21
A	VAN EIJK J H ET AL.: "Nonamylolytic enzymes. In: Baked goods freshness. Technology, evaluation, and inhibition of staling (pp 131-150)" 1996, MARCEL DEKKER, INC., NEW YORK, USA XP002186665 the whole document	21
A	GRUNDY J G: "Preservatives. In: Baked goods freshness. Technology, evaluation, and inhibition of staling (pp 189-203)." 1996, MARCEL DEKKER, INC., NEW YORK, USA XP002186666 page 197, line 15 -page 198, line 30 page 201, line 1-19	25,26
A	POTUS J: "L'intérêt des enzymes en panification. Role of enzymes in breadmaking" COMPTES RENDUS DE L'ACADEMIE D'AGRICULTURE DE FRANCE, vol. 81, no. 2, 1995, pages 27-36, XP000946072 ISSN: 0989-6988 abstract	21

ti d Application No
PUI/IN 01/00094

Patent document cited in search report	Publication date		Patent family member(s)	Publication date
JP 61070980	11-04-1986	JP JP	1384674 C 61052677 B	26-06-1987 14-11-1986
EP 0585988	A 09-03-1994	EP AT AU CA DE DK ES GR JP US	0585988 A1 135163 T 4216693 A 2101308 A1 69301796 D1 69301796 T2 585988 T3 2087646 T3 3019731 T3 6169681 A 6251444 B1	09-03-1994 15-03-1996 03-02-1994 28-01-1994 18-04-1996 01-08-1996 24-06-1996 16-07-1996 31-07-1996 21-06-1994 26-06-2001
EP 0913092	06-05-1999	EP JP	0913092 A2 11192052 A	06-05-1999 21-07-1999
US 4308284	29-12-1981	JP JP JP	1119551 C 55144884 A 57000034 B	28-10-1982 12-11-1980 05-01-1982
EP 1022329	A 26-07-2000	FR AU BR EP WO	2788782 A1 3059800 A 0007600 A 1022329 A1 0043496 A1	28-07-2000 07-08-2000 30-10-2001 26-07-2000 27-07-2000
WO 9928366	10-06-1999	US AU WO US US	5990266 A 1709099 A 9928366 A1 6121033 A 6191176 B1	23-11-1999 16-06-1999 10-06-1999 19-09-2000 20-02-2001

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:
☐ BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
FADED TEXT OR DRAWING
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
☐ GRAY SCALE DOCUMENTS
☐ LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
OTHER:

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.